

EULER'S CONSTANT, q -LOGARITHMS, AND FORMULAS OF RAMANUJAN AND GOSPER

JONATHAN SONDOW (New York) and WADIM ZUDILIN[‡] (Moscow)

26 March 2003

ABSTRACT. The aim of the paper is to relate computational and arithmetic questions about Euler's constant γ with properties of the values of the q -logarithm function, with natural choice of q . By these means, we generalize a classical formula for γ due to Ramanujan, together with Vacca's and Gosper's series for γ , as well as deduce irrationality criteria and tests and new asymptotic formulas for computing Euler's constant. The main tools are Euler-type integrals and hypergeometric series.

We recall the definition of Euler's constant:

$$\begin{aligned}\gamma &= \lim_{n \rightarrow \infty} \left(\sum_{\nu=1}^n \frac{1}{\nu} - \log n \right) \\ &= 0.57721566490153286060651209008240243104215933593992 \dots\end{aligned}$$

and the fact that the (expected) irrationality of γ has not yet been proved.

Recently, the first author gave [9], [10] a construction of $\mathbb{Z}$ -linear forms involving γ and logarithms; namely, he proved that

$$d_{2n} I_n \in \mathbb{Z} + \mathbb{Z}\gamma + \mathbb{Z} \log(n+1) + \mathbb{Z} \log(n+2) + \dots + \mathbb{Z} \log(2n),$$

where

$$\begin{aligned}I_n &= \iint_{[0,1]^2} \frac{x^n(1-x)^n y^n(1-y)^n}{(1-xy)|\log xy|} dx dy \\ &= \sum_{\nu=n+1}^{\infty} \int_{\nu}^{\infty} \left(\frac{n!}{t(t+1) \cdots (t+n)} \right)^2 dt\end{aligned}$$

[‡]The work of the second author is supported by an Alexander von Humboldt research fellowship.

Key words and phrases. Euler's constant, q -logarithm, hypergeometric series, Euler's transform, irrationality.

2000 *Mathematics Subject Classification.* Primary 11Y60; Secondary 11J72, 33C20, 33D15.

and d_n denotes the least common multiple of the numbers $1, 2, \dots, n$. This type of approximation allows deducing several (ir)rationality criteria for Euler's constant [9]. In a sense, what follows is inspired by considerations in [9].

A special function that we will require is the q -logarithm

$$(1) \quad \ln_q(1+z) = \sum_{\nu=1}^{\infty} \frac{(-1)^{\nu-1} z^{\nu}}{q^{\nu} - 1}, \quad |z| < q,$$

for q real with $|q| > 1$ (in fact, our main choices will be $q = 2$ in Sections 2–4 and $q = 3$ in Section 8, but we let $q > 1$ be any integer in Section 9). The function (1) is a q -extension of the ordinary logarithm function

$$\log(1+z) = \sum_{\nu=1}^{\infty} \frac{(-1)^{\nu-1} z^{\nu}}{\nu}, \quad |z| \leq 1, \quad z \neq -1,$$

since

$$\lim_{\substack{q \rightarrow 1 \\ q > 1}} ((q-1) \ln_q(1+z)) = \log(1+z), \quad |z| < 1.$$

We also mention that the irrationality of the values of the q -logarithm (1) for $q \in \mathbb{Z} \setminus \{0, \pm 1\}$ and $z \in \mathbb{Q}$, $z \leq 1$, is known [3].

CONTENTS

1. Gosper's acceleration of Vacca's series.....	2
2. A q -logarithm approach.....	4
3. Irrationality tests for Euler's constant.....	8
4. Computing Euler's constant.....	10
5. Ramanujan's base q integral for Euler's constant.....	11
6. A base q generalization of Gosper's Series.....	12
7. Base q integrals.....	13
8. Base 3 irrationality testing.....	15
9. Base q asymptotic formulas for Euler's constant.....	18
10. Concluding remarks.....	22
References.....	23

1. GOSPER'S ACCELERATION OF VACCA'S SERIES

The following expression for γ is known as Vacca's series [11]:

$$(2) \quad \gamma = \sum_{n=1}^{\infty} (-1)^n \frac{\lfloor \log_2 n \rfloor}{n},$$

where $\log_q x = (\log x)/(\log q)$ is the ordinary logarithm base q . Gosper [6] has transformed this series into a more rapidly converging series of positive rationals. His method (see Section 6) is to use partial summation to write (2) as the double series

$$(3) \quad \gamma = \sum_{\nu=1}^{\infty} \sum_{k=0}^{\infty} \frac{(-1)^k}{2^{\nu} + k},$$

apply Euler's transformation to each inner sum on k , obtaining the double series (7) below, then triangularize it. In this way, he arrives at a version of series (4). Here we give a different proof of (4). To begin, we reverse the order of summation in (3) and write the resulting double series as the integral (5), which is a form of an integral that Bromwich attributes to Catalan ([4], p. 526).

Theorem 1 (Gosper's Series). *We have the series for Euler's constant*

$$(4) \quad \gamma = \frac{1}{2} + \sum_{\nu=2}^{\infty} \frac{1}{2^{\nu+1}} \sum_{k=1}^{\nu-1} \binom{2^{\nu-k} + k}{k}^{-1}.$$

Proof. After expanding $1/(1+x)$ in a geometric series, termwise integration shows that the formula

$$(5) \quad \gamma = \int_0^1 \frac{1}{1+x} \sum_{\nu=1}^{\infty} x^{2^{\nu}-1} dx$$

is equivalent to (3) with the order of summation reversed, which is easily justified by grouping terms in pairs. Integrating by parts $K+1$ times gives

$$(6) \quad \gamma = \sum_{k=0}^K \sum_{\nu=1}^{\infty} \frac{1}{2^{\nu+k+1} \binom{2^{\nu}+k}{k}} + \int_0^1 \frac{(K+1)!}{(1+x)^{K+2}} \sum_{\nu=1}^{\infty} \frac{x^{2^{\nu}+K}}{2^{\nu}(2^{\nu}+1) \cdots (2^{\nu}+K)} dx.$$

Since the integral is less than

$$\int_0^1 \frac{1}{(1+x)^{K+2}} \sum_{\nu=1}^{\infty} \frac{1}{2^{\nu}} dx = \frac{1}{K+1} \left(1 - \frac{1}{2^{K+1}} \right),$$

as K tends to infinity (6) becomes

$$(7) \quad \gamma = \sum_{k=0}^{\infty} \sum_{\nu=1}^{\infty} \frac{1}{2^{\nu+k+1} \binom{2^{\nu}+k}{k}}.$$

The inner series with $k=0$ sums to $1/2$, and in the series with $k>0$ we may collect terms (triangularize) as in (4), completing the proof. $\square$

Remark. We can regard the series on the right of (2), (4), (7) as base $q=2$ series for Euler's constant. Gosper indicated [6] the following extensions of (2), (7) to base $q=3$:

$$(8) \quad \gamma = 2 \sum_{n=1}^{\infty} \frac{[\log_3 n] \cdot \cos \frac{2\pi n}{3}}{n} = 2 \sum_{\nu=1}^{\infty} \sum_{k=0}^{\infty} \frac{\cos \frac{\pi(k-1)}{6}}{3^{\nu+(k+1)/2} \binom{3^{\nu}+k}{k}}$$

(see Sections 5 and 6 for proofs and generalizations), which give faster convergence than formulas (2), (7), respectively.

2. A q -LOGARITHM APPROACH

Lemma 1. *Let q be real, $q > 1$. Then for all integers $n \geq 0$ and k , with $0 < k < q^{n+1}$, the following identity holds:*

$$(9) \quad \sum_{\nu=n+1}^{\infty} \frac{1}{q^{\nu} + k} = \frac{1}{k} \ln_q \left(1 + \frac{k}{q^n} \right).$$

Remark. For $k = 0$, the series in (9) equals

$$\sum_{\nu=n+1}^{\infty} \frac{1}{q^{\nu}} = \frac{q^{-(n+1)}}{1 - q^{-1}} = \frac{1}{q^n(q - 1)}.$$

Proof. We have

$$\begin{aligned} \sum_{\nu=n+1}^{\infty} \frac{1}{q^{\nu} + k} &= \sum_{\nu=n+1}^{\infty} \frac{1}{q^{\nu}} \frac{1}{1 + k \cdot q^{-\nu}} = \sum_{\nu=n+1}^{\infty} \frac{1}{q^{\nu}} \sum_{\mu=1}^{\infty} \frac{(-k)^{\mu-1}}{q^{\nu(\mu-1)}} \\ &= \sum_{\mu=1}^{\infty} (-k)^{\mu-1} \sum_{\nu=n+1}^{\infty} \frac{1}{q^{\nu\mu}} = \sum_{\mu=1}^{\infty} (-k)^{\mu-1} \frac{q^{-(n+1)\mu}}{1 - q^{-\mu}} \\ &= \frac{1}{k} \sum_{\mu=1}^{\infty} \frac{(-1)^{\mu-1} (k/q^n)^{\mu}}{q^{\mu} - 1} = \frac{1}{k} \ln_q \left(1 + \frac{k}{q^n} \right) \end{aligned}$$

as required. $\square$

Remark. P. Sebah [8] has pointed out that in order to compute the q -logarithm, the following formula is much more efficient than (1) or (9):

$$\ln_q(1 + z) = z \sum_{\nu=1}^N \frac{1}{q^{\nu} + z} + r_N(z), \quad \text{where} \quad r_N(z) = \sum_{\nu=1}^{\infty} \frac{(-1)^{\nu-1} z^{\nu}}{q^{N\nu}(q^{\nu} - 1)},$$

with N being any positive integer. The proof of the formula is similar to that of Lemma 1.

Let m be a non-negative integer. Using the rational function

$$(10) \quad R_m(t) = \frac{m!}{t(t+1) \cdots (t+m)} = \sum_{j=0}^m \frac{(-1)^j \binom{m}{j}}{t+j},$$

for each $\nu \geq 0$ define the convergent series

$$(11) \quad S_{\nu,m} = \sum_{t=0}^{\infty} (-1)^t R_m(2^{\nu} + t).$$

Lemma 2. *We have*

$$(12) \quad S_{\nu,m} = 2^m \sum_{k=0}^{\infty} \frac{(-1)^k}{2^{\nu} + k} - \sum_{j=1}^m \binom{m}{j} \sum_{k=0}^{j-1} \frac{(-1)^k}{2^{\nu} + k}.$$

(We use the standard convention that any sum $\sum_{j=a}^b$ is zero if $a > b$.)

Proof. Indeed,

$$\begin{aligned} S_{\nu,m} &= \sum_{j=0}^m \sum_{t=0}^{\infty} \frac{(-1)^{t+j} \binom{m}{j}}{2^{\nu} + t + j} = \sum_{j=0}^m \binom{m}{j} \sum_{t=0}^{\infty} \frac{(-1)^{t+j}}{2^{\nu} + t + j} \\ &= \sum_{j=0}^m \binom{m}{j} \sum_{k=j}^{\infty} \frac{(-1)^k}{2^{\nu} + k} = \sum_{j=0}^m \binom{m}{j} \left(\sum_{k=0}^{\infty} - \sum_{k=0}^{j-1} \right) \frac{(-1)^k}{2^{\nu} + k}, \end{aligned}$$

and (12) follows. $\square$

Lemma 3. *For the series $S_{\nu,m}$, we have the representations*

$$(13) \quad S_{\nu,m} = (-1)^{2^{\nu}} \sum_{t=2^{\nu}}^{\infty} (-1)^t R_m(t) = \int_0^1 \frac{(1-x)^m}{1+x} x^{2^{\nu}-1} dx.$$

Proof. Replace t by $t - 2^{\nu}$ in (11) to get the first equality in (13). For the second, use the Binomial Theorem to write

$$R_m(t) = \int_0^1 (1-x)^m x^{t-1} dx.$$

Substitute this in the series for $S_{\nu,m}$, interchange summation and integration, and sum the series

$$(14) \quad \sum_{t=2^{\nu}}^{\infty} (-1)^{t-2^{\nu}} x^{t-1} = \frac{x^{2^{\nu}-1}}{1+x},$$

arriving at the second equality in (13). To justify termwise integration, replace the geometric series (14) with a finite sum plus remainder $r_N(x) = (-1)^N x^{N+2^{\nu}-1} / (1+x)$, and note that

$$\left| \int_0^1 r_N(x) dx \right| < \frac{1}{N+2^{\nu}} \rightarrow 0 \quad \text{as } N \rightarrow \infty. \quad \square$$

Our final object is the double series

$$(15) \quad I_{n,m} = \sum_{\nu=n+1}^{\infty} S_{\nu,m},$$

where $n \geq 0$. By Lemma 2 and formula (3), for the quantity (15) we have

$$\begin{aligned} I_{n,m} &= 2^m \left(\sum_{\nu=1}^{\infty} - \sum_{\nu=1}^n \right) \sum_{k=0}^{\infty} \frac{(-1)^k}{2^{\nu} + k} - \sum_{j=1}^m \binom{m}{j} \sum_{k=0}^{j-1} \sum_{\nu=n+1}^{\infty} \frac{(-1)^k}{2^{\nu} + k} \\ &= 2^m \gamma - 2^m \sum_{\nu=1}^n \left(\sum_{l=1}^{\infty} - \sum_{l=1}^{2^{\nu}-1} \right) \frac{(-1)^l}{l} - \sum_{j=1}^m \binom{m}{j} \sum_{\nu=n+1}^{\infty} \frac{1}{2^{\nu}} \\ &\quad - \sum_{j=1}^m \binom{m}{j} \sum_{k=1}^{j-1} \sum_{\nu=n+1}^{\infty} \frac{(-1)^k}{2^{\nu} + k}. \end{aligned}$$

For simplicity, assume now that $m \leq 2^{n+1}$. Then using Lemma 1,

$$\begin{aligned} (16) \quad I_{n,m} &= 2^m \cdot (\gamma + n \log 2) - 2^m \sum_{\nu=1}^n \sum_{l=1}^{2^{\nu}-1} \frac{(-1)^{l-1}}{l} - \frac{2^m - 1}{2^n} \\ &\quad - \sum_{j=2}^m \binom{m}{j} \sum_{k=1}^{j-1} \frac{(-1)^k}{k} \ln_2 \left(1 + \frac{k}{2^n} \right). \end{aligned}$$

We can summarize the result of these computations in the following way.

Lemma 4. *If $m \leq 2^{n+1}$, then*

$$(17) \quad I_{n,m} = 2^m \gamma + L_{n,m} - A_{n,m},$$

where

$$L_{n,m} = 2^m n \log 2 + \sum_{j=2}^m \binom{m}{j} \sum_{k=1}^{j-1} \frac{(-1)^{k-1}}{k} \ln_2 \left(1 + \frac{k}{2^n} \right)$$

and

$$A_{n,m} = \frac{2^m - 1}{2^n} + 2^m \sum_{\nu=1}^n \sum_{l=1}^{2^{\nu}-1} \frac{(-1)^{l-1}}{l}$$

satisfy

$$d_m L_{n,m} \in \mathbb{Z} \log 2 + \mathbb{Z} \ln_2 \left(1 + \frac{1}{2^n} \right) + \mathbb{Z} \ln_2 \left(1 + \frac{2}{2^n} \right) + \cdots + \mathbb{Z} \ln_2 \left(1 + \frac{m-1}{2^n} \right)$$

and $d_{2^n} A_{n,m} \in \mathbb{Z}$.

Lemma 5. *For the double series $I_{n,m}$, we have the integral*

$$(18) \quad I_{n,m} = \int_0^1 \frac{(1-x)^m}{1+x} \sum_{\nu=n+1}^{\infty} x^{2^{\nu}-1} dx,$$

which generalizes Catalan's integral (5) for γ .

Proof. Integrating termwise (since the series in the integrand is dominated by a geometric series, the justification in the proof of Lemma 3 applies here) and using Lemma 3, the formula follows from the definition of $I_{n,m}$. The case $I_{0,0} = \gamma$ (see (16)) is indeed the integral (5). $\square$

Lemma 6. For m a positive integer and $r \geq 1$ a real number, we have the bounds

$$(19) \quad \int_0^1 x^{rm-1}(1-x)^m dx > \frac{1}{rm} \left(\frac{r^r}{(r+1)^{r+1}} \right)^m$$

and, for $0 < x < 1$,

$$(20) \quad x^{rm-1}(1-x)^m < 4 \cdot \left(\frac{r^r}{(r+1)^{r+1}} \right)^m.$$

Proof. Euler's beta integral

$$\int_0^1 x^{\alpha-1}(1-x)^{\beta-1} dx = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}, \quad \operatorname{Re} \alpha > 0, \quad \operatorname{Re} \beta > 0,$$

gives

$$\int_0^1 x^{rm-1}(1-x)^m dx = \frac{\Gamma(rm)\Gamma(m+1)}{\Gamma(rm+m+1)} = \frac{1}{rm} \prod_{j=1}^m \left(1 + \frac{rm}{j} \right)^{-1};$$

denote the product by $\Pi(m, r)$. Using $r \geq 1$ to get

$$\frac{j+r(m+1)}{j+rm} < 1 + \frac{1}{m} \leq \left(1 + \frac{1}{r} \right)^{r/m} \quad \text{for } j = 1, 2, \dots, m,$$

we obtain

$$\frac{\Pi(m+1, r)}{\Pi(m, r)} = \frac{1}{r+1} \prod_{j=1}^m \frac{j+rm}{j+r(m+1)} > \frac{1}{r+1} \left(1 + \frac{1}{r} \right)^{-r} = \frac{r^r}{(r+1)^{r+1}},$$

and by induction on m it follows that

$$\Pi(m, r) > \left(\frac{r^r}{(r+1)^{r+1}} \right)^m$$

since the inequality holds for $m = 1$. This proves (19).

If $0 < x < 1$, then

$$\begin{aligned} x^{rm-1}(1-x)^m &< (x^r(1-x))^{m-1/r} \leq \left(\frac{r^r}{(r+1)^{r+1}} \right)^{m-1/r} \\ &= \frac{(r+1)^{1+1/r}}{r} \cdot \left(\frac{r^r}{(r+1)^{r+1}} \right)^m \end{aligned}$$

and (20) follows, using $r \geq 1$. $\square$

Lemma 7. *If $n \geq 0$ is an integer and r is a real number satisfying $1 \leq r \leq 2^{n+1}$, then we have the bounds*

$$(21) \quad \frac{1}{2r(m+1)} \left(\frac{r^r}{(r+1)^{r+1}} \right)^{m+1} < I_{n,m} < 6 \cdot \left(\frac{r^r}{(r+1)^{r+1}} \right)^m \quad \text{for } m = \left\lfloor \frac{2^{n+1}}{r} \right\rfloor.$$

In particular, for $r \geq 1$ fixed and $m = \lfloor 2^{n+1}/r \rfloor \geq 1$, the quantity $I_{n,m}$ is positive and decreases exponentially at the rate

$$(22) \quad \lim_{n \rightarrow \infty} \frac{\log I_{n, \lfloor 2^{n+1}/r \rfloor}}{\lfloor 2^{n+1}/r \rfloor} = r \log r - (r+1) \log(r+1).$$

Proof. Put $m = \lfloor 2^{n+1}/r \rfloor$ in Lemma 5. Then $m+1 > 2^{n+1}/r$, hence

$$I_{n,m} > \int_0^1 \frac{(1-x)^m}{1+x} x^{2^{n+1}-1} dx > \frac{1}{2} \int_0^1 x^{r(m+1)-1} (1-x)^{m+1} dx,$$

and the lower bound in (21) follows from (19) (with $m+1$ in place of m).

Using $m \leq 2^{n+1}/r$ and relations (20) and (5), we have

$$\begin{aligned} I_{n,m} &\leq \int_0^1 \frac{(1-x)^m}{1+x} x^{rm-1} \sum_{\nu=n+1}^{\infty} x^{2^\nu-2^{n+1}} dx \\ &< \int_0^1 x^{rm-1} (1-x)^m \frac{1}{1+x} \left(1 + \sum_{\nu=1}^{\infty} x^{2^\nu-1} \right) dx \\ &< 4 \left(\frac{r^r}{(r+1)^{r+1}} \right)^m (\log 2 + \gamma), \end{aligned}$$

and the upper bound in (21) follows from the inequality $4(\log 2 + \gamma) < 6$. Since (21) implies (22), we are done. $\square$

Remark 1. If $2^{n+1}/r$ is an integer, then we may replace $m+1$ by m in the lower bound in (21).

Remark 2. As $r \rightarrow \infty$, the right-hand side of (22) equals $-\log r + O(1)$, hence its absolute value tends to infinity with r .

3. IRRATIONALITY TESTS FOR EULER'S CONSTANT

Theorem 2 (Rationality Criterion for γ). *The fractional part of $d_{2^n} L_{n,m}$ equals $d_{2^n} I_{n,m}$ for all n and m with*

$$(23) \quad n \text{ sufficiently large and } 2^{n-1} \leq m \leq 2^{n+1},$$

if and only if Euler's constant is a rational number.

Proof. With $2^{n-1} \leq m \leq 2^{n+1}$ set $r = 2^{n+1}/m$, so that $1 \leq r \leq 4$. By Lemma 7, we have $I_{n,m} < 6\rho(r)^m = 6\rho(r)^{2^n \cdot 2/r}$, where $\rho(r) = r^r/(r+1)^{r+1}$. Straightforward verification shows that $\rho(r)^{2/r}$ is increasing for $r > 0$, and satisfies

$$0 < \rho(r)^{2/r} < \frac{1}{e} \quad \text{for } 0 < r < r_0 = 5.6213305349\dots,$$

where $\rho(r_0)^{2/r_0} = 1/e$. Therefore $I_{n,m} < 6(\rho(4)^{1/2})^{2^n}$ and $\rho(4)^{1/2} < 1/e$. Since the Prime Number Theorem implies that for any $\varepsilon > 0$ we have $d_N = O((e^{1+\varepsilon})^N)$ as $N \rightarrow \infty$, we conclude that $0 < d_{2^n} I_{n,m} < 1$ for n and m satisfying (23).

Now multiply (17) by d_{2^n} and write the result as

$$(24) \quad d_{2^n} L_{n,m} = d_{2^n} I_{n,m} + d_{2^n} (A_{n,m} - 2^m \gamma).$$

Since $d_{2^n} A_{n,m} \in \mathbb{Z}$ by Lemma 4, the theorem follows. $\square$

Remark. The argument shows that the Criterion holds with (23) replaced by the more general conditions $n \geq n(r_1)$ and $\lceil 2^{n+1}/r_1 \rceil \leq m \leq 2^{n+1}$, where r_1 is any number between 1 and r_0 .

Theorem 3 (Irrationality Test for γ). *If the fractional part of $d_{2^n} L_{n,2^n}$ satisfies*

$$(25) \quad \{d_{2^n} L_{n,2^n}\} > 6 \cdot \left(\frac{128}{729}\right)^{2^{n-1}}$$

infinitely often, then Euler's constant is irrational. In fact, the inequality for a given $n > 0$ implies that if γ is a rational number, then its denominator does not divide $2^{2^n} d_{2^n}$.

Proof. It follows from [7], Theorem 13, that $d_{2N} < 8^N$ for all integers $N \geq 1$ (see [9], Lemma 3). Combined with the upper bound in Lemma 7 for $r = 2$, this gives

$$0 < d_{2^n} I_{n,2^n} < 8^{2^{n-1}} \cdot 6 \cdot \left(\frac{4}{27}\right)^{2 \cdot 2^{n-1}} = 6 \cdot \left(\frac{128}{729}\right)^{2^{n-1}}$$

for $n \geq 1$. The first part of the theorem follows by the Rationality Criterion for γ , and the second part by formula (24) with $m = 2^n$. $\square$

Remark 1. Condition (25) in Theorem 3 can be replaced by the simpler condition

$$\{d_{2^n} L_{n,2^n}\} > 5^{-2^{n-1}}$$

if $n > 1$. For $n > 4$, this follows from the inequalities $128/729 = 0.17558299\dots < 1/5$ and $6 \cdot (128/729)^{14} < 5^{-14}$. For $n = 2, 3, 4$, it follows from the proof of Theorem 3 by

replacing the bound $d_{2m} < 8^m$ with the exact values $d_4 = 12$, $d_8 = 840$, $d_{16} = 720720$, for which $d_{2^n} \cdot 6 \cdot (4/27)^{2^n} < 5^{-2^{n-1}}$.

Remark 2. Using the Rationality Criterion for γ , we may generalize the Irrationality Test as follows: γ is irrational if there exist infinitely many integers n and m such that $2^{n-1} \leq m \leq 2^{n+1}$ and

$$(26) \quad \{d_{2^n} L_{n,m}\} > 8^{2^{n-1}} \cdot 6 \cdot \left(\frac{r^r}{(r+1)^{r+1}} \right)^m, \quad \text{where } r = \frac{2^{n+1}}{m}.$$

For example, the case $m = 2^n$ is Theorem 3, and the case $m = 2^{n+1}$ can be stated: a sufficient condition for irrationality of γ is that

$$\{d_{2^n} L_{n,2^{n+1}}\} > 6 \cdot 2^{-5 \cdot 2^{n-1}} \quad \text{infinitely often.}$$

For any $\varepsilon > 0$, we may further refine the Test by replacing $8^{2^{n-1}} \cdot 6$ with $e^{(1+\varepsilon)2^n}$ in (26).

Example. P. Sebah [8] has calculated that

$$\{d_{2^{12}} L_{12,2^{12}}\} = 0.178346164 \dots$$

Since the last number exceeds $5^{-2^{11}}$, if γ is a rational number a/b , then b is not a divisor of $2^{4096} \cdot d_{4096}$; in particular, $|b| \geq 4099$ (since the numbers $4097 = 17 \cdot 241$ and $4098 = 2 \cdot 3 \cdot 683$ divide d_{4096}).

Remark. Similar irrationality tests for γ , but using ordinary logarithms instead of 2-logarithms, are proved in [9], where an example shows that no divisor of $\binom{20000}{10000} \cdot d_{20000}$ can be a denominator of γ . The present example yields additional information because of the high power of 2.

4. COMPUTING EULER'S CONSTANT

Theorem 4. The following asymptotic formulas as $n \rightarrow \infty$ are valid:

$$(27) \quad \gamma = \frac{A_{n,2^{n+1}} - L_{n,2^{n+1}}}{2^{2^{n+1}}} + O(2^{-6 \cdot 2^n}),$$

$$(28) \quad \gamma = \frac{A_{n,2^n} - L_{n,2^n}}{2^{2^n}} + O\left(\left(\frac{2}{27}\right)^{2^n}\right),$$

$$(29) \quad \gamma = \frac{A_{n,2^{n-1}} - L_{n,2^{n-1}}}{2^{2^{n-1}}} + O\left(\left(\frac{128}{3125}\right)^{2^{n-1}}\right),$$

$$(30) \quad \gamma = \frac{1}{2} A_{n,1} - n \log 2 + O(2^{-n}).$$

Proof. For (27), put $m = 2^{n+1}$ in (17) and use the upper bound in (21) with $r = 1$. For (28), (29), take $m = 2^n, 2^{n-1}$ and $r = 2, 4$, respectively. Finally, let $m = 1$ in Lemma 4 to get $I_{n,1} = 2\gamma + 2n \log 2 - A_{n,1}$. From (21) with $r = 2^{n+1}$, we obtain

$$0 < I_{n,1} < 6 \cdot \frac{(2^{n+1})^{2^{n+1}}}{(2^{n+1} + 1)^{2^{n+1}+1}} = \frac{6}{2^{n+1} + 1} \left(1 + \frac{1}{2^{n+1}}\right)^{-2^{n+1}} < 2^{-n+1}$$

for $n \geq 0$, proving more than (30). $\square$

Remark. Using (28) and his calculation of $L_{n,2^n}$ for $n = 12$, P. Sebah [8] has computed γ correct to 4631 decimal places, which is the accuracy $(2/27)^{2^{12}}$ predicted. Formula (27) gives almost 60% more digits of γ , but requires computing about twice as many 2-logarithms. Similarly, (29) yields nearly 39% fewer digits than (28), but involves only half as many 2-logarithms. Letting m continue down to 1, we reach formula (30), which doesn't involve any 2-logarithms.

5. RAMANUJAN'S BASE q INTEGRAL FOR EULER'S CONSTANT

For integer $q \geq 2$, Ramanujan (see [2] for proofs and references) gave the formula for Euler's constant

$$(31) \quad \gamma = \int_0^1 \left(\frac{q}{1-x^q} - \frac{1}{1-x} \right) \sum_{\nu=1}^{\infty} x^{q^{\nu}-1} dx,$$

which reduces to Catalan's integral (5) when $q = 2$. Equation (31) implies the following base q generalization of Vacca's series (2) (see [2], Theorem 2.6):

$$(32) \quad \gamma = \sum_{n=1}^{\infty} \frac{\sigma_n \lfloor \log_q n \rfloor}{n},$$

where

$$(33) \quad \sigma_n = \sigma_{n,q} = \begin{cases} q-1 & \text{if } q \mid n, \\ -1 & \text{if } q \nmid n. \end{cases}$$

The case $q = 3$ gives the first equality in (8).

Let $\epsilon = \epsilon_q$ be a fixed primitive q -th root of unity (for instance, $\epsilon = e^{2\pi i/q}$). Using

$$\sigma_n = -1 + \sum_{l=0}^{q-1} \epsilon^{nl} = \sum_{l=1}^{q-1} \epsilon^{nl},$$

we can represent formula (32) as

$$(34) \quad \gamma = \sum_{l=1}^{q-1} \sum_{n=1}^{\infty} \frac{\lfloor \log_q n \rfloor \epsilon^{nl}}{n},$$

which can be regarded as another base q generalization of Vacca's series (2).

6. A BASE q GENERALIZATION OF GOSPER'S SERIES

Following Gosper's proof [6] of (7), we write (34) as

$$\gamma = \sum_{l=1}^{q-1} \sum_{\nu=1}^{\infty} \sum_{k=0}^{\infty} \frac{\epsilon^{kl}}{q^{\nu} + k} = \sum_{l=1}^{q-1} \sum_{\nu=1}^{\infty} \frac{1}{q^{\nu}} \cdot {}_2F_1\left(\begin{matrix} 1, q^{\nu} \\ q^{\nu} + 1 \end{matrix} \middle| \epsilon^l\right)$$

and apply Euler's transform (see, e.g., [1], Section 2.4, formula (1))

$${}_2F_1\left(\begin{matrix} \alpha, \beta \\ \gamma \end{matrix} \middle| z\right) = \frac{1}{(1-z)^{\alpha}} \cdot {}_2F_1\left(\begin{matrix} \alpha, \gamma - \beta \\ \gamma \end{matrix} \middle| \frac{-z}{1-z}\right),$$

yielding

$$\begin{aligned} \gamma &= \sum_{l=1}^{q-1} \sum_{\nu=1}^{\infty} \frac{1}{q^{\nu}} \cdot \frac{1}{1 - \epsilon^l} \cdot {}_2F_1\left(\begin{matrix} 1, 1 \\ q^{\nu} + 1 \end{matrix} \middle| \frac{-\epsilon^l}{1 - \epsilon^l}\right) \\ &= \sum_{l=1}^{q-1} \sum_{\nu=1}^{\infty} \frac{1}{q^{\nu}} \sum_{k=0}^{\infty} \binom{q^{\nu} + k}{k}^{-1} \frac{(-1)^k \epsilon^{kl}}{(1 - \epsilon^l)^{k+1}}. \end{aligned}$$

Substituting $x = 1$ into the expansion

$$\prod_{j=1}^{q-1} (1 - \epsilon^j x) = \frac{1 - x^q}{1 - x} = 1 + x + x^2 + \cdots + x^{q-1}$$

leads to

$$(35) \quad \frac{1}{1 - \epsilon^l} = \frac{1}{q} \cdot \prod_{\substack{j=1 \\ j \neq l}}^{q-1} (1 - \epsilon^j)$$

(where the empty product in the case $q = 2$ equals 1). Therefore, we arrive at the following result.

Theorem 5. *We have the base $q \geq 2$ accelerated series*

$$(36) \quad \gamma = \sum_{\nu=1}^{\infty} \sum_{k=0}^{\infty} \frac{(-1)^k \chi_q(k)}{q^{\nu+k+1} \binom{q^{\nu} + k}{k}},$$

where

$$\chi_q(k) = \sum_{l=1}^{q-1} \epsilon^{kl} \prod_{\substack{j=1 \\ j \neq l}}^{q-1} (1 - \epsilon^j)^{k+1},$$

a symmetric polynomial in the roots of the polynomial $1 + x + x^2 + \cdots + x^q$, is an integer-valued function independent of the choice of primitive q -th root of unity ϵ .

Remark. From (35) we have

$$\epsilon^{kl} \prod_{\substack{j=1 \\ j \neq l}}^{q-1} (1 - \epsilon^j)^{k+1} = \epsilon^{kl} \frac{q^{k+1}}{(1 - \epsilon^l)^{k+1}},$$

and it follows that

$$|\chi_q(k)| \leq (q-1) \cdot \frac{q^{k+1}}{|1 - e^{2\pi i/q}|^{k+1}} = (q-1) \left(\frac{q}{2 \sin \frac{\pi}{q}} \right)^{k+1}.$$

Thus we have the relation

$$\limsup_{k \rightarrow \infty} |\chi_q(k)|^{1/k} \leq \frac{q}{2 \sin \frac{\pi}{q}},$$

which shows how the convergence of series (34) has been accelerated in (36).

Examples. Series (36) is a generalization of Gosper's series (7), which is the case $q = 2$. When $q = 3$, taking $\epsilon = e^{2\pi i/3}$ we obtain

$$\begin{aligned} \chi_3(k) &= e^{2\pi ki/3} (1 - e^{-2\pi i/3})^{k+1} + e^{-2\pi ki/3} (1 - e^{2\pi i/3})^{k+1} \\ &= e^{2\pi ki/3} \cdot 3^{(k+1)/2} e^{\pi(k+1)i/6} + e^{-2\pi ki/3} \cdot 3^{(k+1)/2} e^{-\pi(k+1)i/6} \\ &= 3^{(k+1)/2} \cdot 2 \cos \frac{\pi(k-1)}{6}, \end{aligned}$$

which proves the second equality in (8). When $q = 4$, we take $\epsilon = e^{\pi i/2} = i$ and obtain

$$\begin{aligned} \chi_4(k) &= i^k 2^{k+1} (1+i)^{k+1} + (-1)^k (1-i)^{k+1} (1+i)^{k+1} + (-i)^k 2^{k+1} (1-i)^{k+1} \\ &= 2^{3(k+1)/2+1} \cdot \cos \frac{\pi(3k+1)}{4} + (-1)^k 2^{k+1}, \end{aligned}$$

which is an integer-valued function of k .

7. BASE q INTEGRALS

There are several ways to generalize integrals (18) for $I_{n,m}$ and (31) for γ simultaneously. One way is to define

$$(37) \quad I_{n,m,q} = \int_0^1 \left(\frac{q}{1-x^q} - \frac{1}{1-x} \right) (1-x)^m \sum_{\nu=n+1}^{\infty} x^{q^\nu-1} dx.$$

Then $I_{n,m,2} = I_{n,m}$ and $I_{0,0,q} = \gamma$, and the analog of Lemma 7 with $r = q$ and $m = q^n$,

$$(38) \quad \lim_{n \rightarrow \infty} \frac{\log I_{n,q^n,q}}{q^n} = q \log q - (q+1) \log(q+1),$$

can be easily derived by the methods of Section 2 (cf. Lemma 9 below). We may expand integral (37) as in the proof of Lemma 4. Namely, with $R_m(t)$ defined in (10), and σ_t in (33), we have

$$\frac{q}{1-x^q} - \frac{1}{1-x} = \frac{(q-1) - x - x^2 - \dots - x^{q-1}}{1-x^q} = \sum_{t=0}^{\infty} \sigma_t x^t,$$

hence

$$\begin{aligned} I_{n,m,q} &= \sum_{t=0}^{\infty} \sigma_t \sum_{\nu=n+1}^{\infty} \int_0^1 x^{q^\nu+t-1} (1-x)^m dx = \sum_{\nu=n+1}^{\infty} \sum_{t=0}^{\infty} \sigma_t R_t(q^\nu + t) \\ &= \sum_{j=0}^m (-1)^j \binom{m}{j} \sum_{\nu=n+1}^{\infty} \sum_{k=j}^{\infty} \frac{\sigma_{k-j}}{q^\nu + k}. \end{aligned}$$

Unfortunately, the last representation shows that $I_{n,m,q}$ involves not only logarithms and q -logarithms but also “generalized Euler constants”

$$(39) \quad \gamma_{j,q} = \sum_{\nu=1}^{\infty} \sum_{k=j}^{\infty} \frac{\sigma_{k-j}}{q^\nu + k}, \quad j = 0, 1, \dots, q-1;$$

the equality $\gamma_{0,q} = \gamma$ follows from (32) just as (3) follows from Vacca’s series (2) (the case $q = 2$). We may avoid the appearance of generalized Euler constants $\gamma_{j,q} \neq \gamma$ in the particular case $q = 3$ (details are in Section 8), but not for $q > 3$ (see Section 10).

However, there is a way to generalize integral (18) for $I_{n,m}$ that solves this problem, namely, by defining

$$\begin{aligned} (40) \quad I'_{n,m,q} &= \int_0^1 \left(\frac{q}{1-x^q} - \frac{1}{1-x} \right) \left(q - \frac{1-x^q}{1-x} \right)^m \sum_{\nu=n+1}^{\infty} x^{q^\nu-1} dx \\ &= \int_0^1 \frac{(q-1-x-x^2-\dots-x^{q-1})^{m+1}}{1-x^q} \sum_{\nu=n+1}^{\infty} x^{q^\nu-1} dx, \end{aligned}$$

with the properties $I'_{n,m,2} = I_{n,m}$ and $I'_{0,0,q} = \gamma$, as for (37). For integral $I'_{n,m,q}$ in (40), we may generalize the results of Sections 2 and 4, but the poor asymptotics of the integral make these generalizations useless for irrationality testing. For example, the choice $m = (q^n - 1)/(q - 1)$ gives the inclusions

$$\begin{aligned} d_{q^n} I'_{n,m,q} &\in \mathbb{Z} + \mathbb{Z}\gamma + \mathbb{Z}\log q + \mathbb{Z}\ln_q \left(1 + \frac{1}{q^n} \right) + \mathbb{Z}\ln_q \left(1 + \frac{2}{q^n} \right) + \dots \\ &\quad + \mathbb{Z}\ln_q \left(1 + \frac{q^n - 2}{q^n} \right) \end{aligned}$$

(see Lemma 10 below) and the asymptotics

$$\lim_{n \rightarrow \infty} \frac{\log I'_{n,(q^n-1)/(q-1),q}}{q^n} > -1$$

(see Lemma 13); this makes it impossible to use $I'_{n,m,q}$ to give irrationality tests for $q > 2$ similar to those of Theorems 2 and 3 for $q = 2$. Instead we use $I'_{n,m,q}$ to extend Theorem 4 to a base $q > 2$ asymptotic formula for γ ; details are in Section 9.

8. BASE 3 IRRATIONALITY TESTING

Taking $q = 3$ in (31) gives one of Ramanujan's formulas for Euler's constant

$$(41) \quad \gamma = \int_0^1 \frac{2+x}{1+x+x^2} \sum_{\nu=1}^{\infty} x^{3^\nu-1} dx$$

(see [2], Corollary 2.3 and the equation following (2.7)), and taking $q = 3$ in (37) gives the base 3 integral

$$(42) \quad I_{n,m,3} = \int_0^1 \frac{2+x}{1+x+x^2} (1-x)^m \sum_{\nu=n+1}^{\infty} x^{3^\nu-1} dx.$$

It turns out that, for m a multiple of 6, we can extend the results of Sections 2–4 to base $q = 3$, as follows.

Since

$$\frac{(1-x)^6}{1+x+x^2} = 28 - 34x + 21x^2 - 7x^3 + x^4 - \frac{27}{1+x+x^2},$$

by induction we have

$$(43) \quad \frac{(1-x)^{6k}}{1+x+x^2} = Q_k(x) + \frac{(-3)^{3k}}{1+x+x^2},$$

where $Q_k(x)$ is the polynomial defined by the recursion

$$(44) \quad \begin{aligned} Q_0(x) &= 0, \\ Q_{k+1}(x) &= (1-x)^6 Q_k(x) + (-3)^{3k} (28 - 34x + 21x^2 - 7x^3 + x^4) \quad \text{for } k \geq 0. \end{aligned}$$

From (42), (43), (41), for $m = 6k$ we obtain

$$(45) \quad \begin{aligned} I_{n,m,3} &= \int_0^1 (2+x) \left(Q_k(x) + \frac{(-3)^{3k}}{1+x+x^2} \right) \left(\sum_{\nu=1}^{\infty} - \sum_{\nu=1}^n \right) x^{3^\nu-1} dx \\ &= (-3)^{3k} \left(\gamma - \int_0^1 \frac{2+x}{1+x+x^2} \sum_{\nu=1}^n x^{3^\nu-1} dx \right) \\ &\quad + \int_0^1 (2+x) Q_k(x) \sum_{\nu=n+1}^{\infty} x^{3^\nu-1} dx. \end{aligned}$$

By division, we find that

$$\frac{(2+x)x^{3^\nu-1}}{1+x+x^2} = 1+x - \sum_{\mu=1}^{3^{\nu-1}-1} (2x^{3^\mu-1} - x^{3^\mu} - x^{3^\mu+1}) - \frac{1+2x}{1+x+x^2}$$

for $\nu \geq 1$, where the sum is zero if $\nu = 1$. Therefore,

$$(46) \quad \int_0^1 \frac{2+x}{1+x+x^2} \sum_{\nu=1}^n x^{3^\nu-1} = n \left(\frac{3}{2} - \log 3 \right) - \sum_{\nu=2}^n \sum_{\mu=1}^{3^{\nu-1}-1} \left(\frac{2}{3^\mu} - \frac{1}{3^\mu+1} - \frac{1}{3^\mu+2} \right)$$

for $n \geq 1$, where the double sum vanishes if $n = 1$.

To evaluate the last integral in (45), assume that $m = 6k < 3^{n+1}$ and let $a_{0,k}, \dots, a_{6k-1,k} \in \mathbb{Z}$ denote the coefficients in the polynomial

$$(47) \quad \sum_{j=0}^{6k-1} a_{j,k} x^j = (2+x)Q_k(x).$$

Noting that $a_{0,k} = (-3)^{3k} 56k$ by (44), and using Lemma 1, we have

$$(48) \quad \begin{aligned} \int_0^1 (2+x)Q_k(x) \sum_{\nu=n+1}^{\infty} x^{3^\nu-1} dx &= \sum_{j=0}^{6k-1} a_{j,k} \sum_{\nu=n+1}^{\infty} \frac{1}{3^\nu + j} \\ &= (-1)^k \frac{28k}{3^{n-3k}} + \sum_{j=1}^{6k-1} \frac{a_{j,k}}{j} \ln_3 \left(1 + \frac{j}{3^n} \right). \end{aligned}$$

Formulas (45), (46), (48) imply the following base 3 version of Lemma 4.

Lemma 8. *If $m < 3^{n+1}$ is a multiple of 6, say $m = 6k$, then*

$$(49) \quad I_{n,m,3} = (-3)^{3k} \gamma + L_{n,m,3} - A_{n,m,3},$$

where

$$L_{n,m,3} = (-3)^{3k} n \log 3 + \sum_{j=1}^{m-1} \frac{a_{j,k}}{j} \ln_3 \left(1 + \frac{j}{3^n} \right)$$

and

$$A_{n,m,3} = (-3)^{3k} \left(\frac{3n}{2} - \frac{28k}{3^n} - \sum_{\nu=2}^n \sum_{\mu=1}^{3^{\nu-1}-1} \left(\frac{2}{3\mu} - \frac{1}{3\mu+1} - \frac{1}{3\mu+2} \right) \right)$$

satisfy

$$d_m L_{n,m,3} \in \mathbb{Z} \log 3 + \mathbb{Z} \ln_3 \left(1 + \frac{1}{3^n} \right) + \mathbb{Z} \ln_3 \left(1 + \frac{2}{3^n} \right) + \dots + \mathbb{Z} \ln_3 \left(1 + \frac{m-1}{3^n} \right)$$

and $d_{3^n} A_{n,m,3} \in \mathbb{Z}$, and the $a_{j,k}$ are integers determined by (47) and (44).

Lemma 9. *We have the bounds*

$$(50) \quad \frac{1}{40m+90} \left(\frac{3^3}{4^4} \right)^m < I_{n,m,3} < 3 \cdot \left(\frac{3^3}{4^4} \right)^m \quad \text{for } m = 3^n - 3 \geq 0$$

and the limit

$$(51) \quad \lim_{n \rightarrow \infty} \frac{\log I_{n,3^n-3,3}}{3^n} = 3 \log 3 - 4 \log 4 = -2.24934057 \dots$$

Proof. Following the proof of Lemma 7, with $m = 3^n - 3$ we have

$$\begin{aligned} I_{n,m,3} &> \int_0^1 \frac{2+x}{1+x+x^2} (1-x)^m x^{3^{n+1}-1} dx > \int_0^1 x^{3m+8} (1-x)^m dx \\ &= \frac{(3m+8)! m!}{(4m+9)!} > \frac{(3/4)^8}{4m+9} \cdot \binom{4m}{m}^{-1} > \frac{1}{40m+90} \left(\frac{3^3}{4^4}\right)^m \end{aligned}$$

and

$$\begin{aligned} I_{n,m,3} &= \int_0^1 \frac{2+x}{1+x+x^2} (1-x)^m x^{3m+8} \sum_{\nu=n+1}^{\infty} x^{3^\nu-3^{n+1}} dx \\ &< \int_0^1 x^{3m+8} (1-x)^m \frac{2+x}{1+x+x^2} \left(1 + \sum_{\nu=1}^{\infty} x^{3^\nu-1}\right) dx \\ &< \left(\frac{3^3}{4^4}\right)^m (2+\gamma) < 3 \cdot \left(\frac{3^3}{4^4}\right)^m. \end{aligned}$$

This proves the lemma. $\square$

Theorem 6. *We have the asymptotic formula for Euler's constant*

$$\gamma = (-1)^{n-1} \frac{A_{n,3^n-3,3} - L_{n,3^n-3,3}}{3^{(3^n-3)/2}} + O\left(\left(\frac{3^{5/2}}{64}\right)^{3^n}\right) \quad \text{as } n \rightarrow \infty.$$

Proof. Let $m = 6k = 3^n - 3$ for $n > 0$. Then $(-1)^{3k} = (-1)^{n-1}$ and the formula follows from (49) and (51). $\square$

Theorem 7. *Euler's constant is rational if and only if the fractional part of $d_{3^n} L_{n,3^n-3,3}$ equals $d_{3^n} I_{n,3^n-3,3}$ for all large n .*

Proof. Multiply (49) by d_{3^n} and write the result as

$$d_{3^n} L_{n,m,3} = d_{3^n} I_{n,m,3} + d_{3^n} (A_{n,m,3} - (-3)^{3k} \gamma).$$

By the Prime Number Theorem and Lemma 9, for $m = 3^n - 3$ and any $\varepsilon > 0$ we have $d_{3^n} I_{n,m,3} = O((3^3 e^{1+\varepsilon}/4^4)^m)$ as $m \rightarrow \infty$. Thus $0 < d_{3^n} I_{n,3^n-3,3} < 1$ for all large n . Since $d_{3^n} A_{n,3^n-3,3} \in \mathbb{Z}$, the theorem follows. $\square$

Theorem 8. *If the inequality*

$$\{d_{3^n} L_{n,3^n-3,3}\} > 3 \cdot \left(\frac{3}{4}\right)^{3^{n+1}}$$

holds infinitely often, then γ is irrational. In fact, the inequality for a given $n > 0$ implies that no divisor of $3^{3^n} d_{3^n}$ can be a denominator of γ .

Proof. The inequalities $d_{3N} \leq d_{4N} < 8^{2N}$ and the upper bound in (50) imply that

$$d_{3^n} I_{n,3^n-3,3} < 8^{2 \cdot 3^{n-1}} \cdot 3 \cdot \left(\frac{3^3}{4^4}\right)^{3^n} = 3 \cdot \left(\frac{3}{4}\right)^{3^{n+1}}$$

for $n \geq 1$. Using Lemma 8, this proves the second part of the theorem, which implies the first. $\square$

9. BASE q ASYMPTOTIC FORMULAS FOR EULER'S CONSTANT

Denoting by $F_q(x)$, $G_q(x)$ the polynomials

$$(52) \quad \begin{aligned} F_q(x) &= 1 + x + x^2 + \cdots + x^{q-1} = \frac{1 - x^q}{1 - x}, \\ G_q(x) &= (q-1) + (q-2)x + \cdots + x^{q-2} = \frac{q - F_q(x)}{1 - x}, \end{aligned}$$

we may write the integral (40) as

$$(53) \quad I'_{n,m,q} = \int_0^1 \frac{G_q(x)}{F_q(x)} (q - F_q(x))^m \left(\sum_{\nu=1}^{\infty} - \sum_{\nu=1}^n \right) x^{q^\nu-1} dx.$$

Expanding the binomial and using Ramanujan's formula $I_{0,0,q} = \gamma$, we obtain

$$(54) \quad \begin{aligned} I'_{n,m,q} &= q^m \left(\gamma - \int_0^1 \frac{G_q(x)}{F_q(x)} \sum_{\nu=1}^n x^{q^\nu-1} dx \right) \\ &\quad + \int_0^1 G_q(x) \sum_{j=1}^m (-1)^j \binom{m}{j} q^{m-j} F_q(x)^{j-1} \sum_{\nu=n+1}^{\infty} x^{q^\nu-1} dx. \end{aligned}$$

In the first integrand, we find by dividing that

$$\frac{G_q(x)x^{q^\nu-1}}{F_q(x)} = F_{q-1}(x) + \sum_{\mu=1}^{q^\nu-1-1} (F_q(x) - q)x^{q^\mu-1} - \frac{F'_q(x)}{F_q(x)}$$

for $\nu \geq 1$, the summation being zero if $\nu = 1$. Therefore,

$$(55) \quad \int_0^1 \frac{G_q(x)}{F_q(x)} \sum_{\nu=1}^n x^{q^\nu-1} dx = n \left(\sum_{\mu=1}^{q-1} \frac{1}{\mu} - \log q \right) - \sum_{\nu=2}^n \sum_{\mu=1}^{q^{\nu-1}-1} \left(\frac{q-1}{q\mu} - \sum_{\lambda=1}^{q-1} \frac{1}{q\mu + \lambda} \right)$$

for $n \geq 1$, where the double summation vanishes if $n = 1$.

In the second integrand in (54), let $b_{0,m,q}, \dots, b_{m(q-1)-1,m,q} \in \mathbb{Z}$ denote the coefficients of the polynomial

$$(56) \quad \sum_{l=0}^{m(q-1)-1} b_{l,m,q} x^l = G_q(x) \sum_{j=1}^m (-1)^j \binom{m}{j} q^{m-j} F_q(x)^{j-1}.$$

Note that $b_{0,m,q} = (q-1)((q-1)^m - q^m)$ by (52). If $m \leq q^{n+1}/(q-1)$, then by Lemma 1 the last integral in (54) equals

$$(57) \quad \begin{aligned} \int_0^1 \sum_{l=0}^{m(q-1)-1} b_{l,m,q} x^l \sum_{\nu=n+1}^{\infty} x^{q^\nu-1} dx &= \sum_{l=0}^{m(q-1)-1} b_{l,m,q} \sum_{\nu=n+1}^{\infty} \frac{1}{q^\nu + l} \\ &= \frac{(q-1)^m - q^m}{q^n} + \sum_{l=1}^{m(q-1)-1} \frac{b_{l,m,q}}{l} \ln_q \left(1 + \frac{l}{q^n} \right). \end{aligned}$$

Equations (54)–(57) imply the following generalization of Lemma 4 to base q .

Lemma 10. *If $m \leq q^{n+1}/(q-1)$, then*

$$(58) \quad I'_{n,m,q} = q^m \gamma + L'_{n,m,q} - A'_{n,m,q},$$

where

$$L'_{n,m,q} = q^m n \log q + \sum_{l=1}^{m(q-1)-1} \frac{b_{l,m,q}}{l} \ln_q \left(1 + \frac{l}{q^n} \right)$$

and

$$A'_{n,m,q} = \frac{q^m - (q-1)^m}{q^n} + q^m \left(\sum_{\mu=1}^{q-1} \frac{n}{\mu} - \sum_{\nu=2}^n \sum_{\mu=1}^{q^{\nu-1}-1} \left(\frac{q-1}{q\mu} - \sum_{\lambda=1}^{q-1} \frac{1}{q\mu + \lambda} \right) \right)$$

satisfy

$$\begin{aligned} d_{m(q-1)} L'_{n,m,q} &\in \mathbb{Z} \log q + \mathbb{Z} \ln_q \left(1 + \frac{1}{q^n} \right) + \mathbb{Z} \ln_q \left(1 + \frac{2}{q^n} \right) + \cdots \\ &\quad + \mathbb{Z} \ln_q \left(1 + \frac{m(q-1)-1}{q^n} \right) \end{aligned}$$

and $d_{q^n} A'_{n,m,q} \in \mathbb{Z}$, and the $b_{l,m,q}$ are integers determined by (56) and (52).

To optimize the asymptotics of $I'_{n,m,q}/q^m$ in the range of Lemma 10, and thus obtain the smallest error term in our formula for γ , we choose $m = (q^n - 1)/(q - 1)$ in the remainder of this section. Let $f_q(x)$ denote the polynomial

$$f_q(x) = (q - F_q(x))x^{q(q-1)} = (q-1)x^{q(q-1)} - \sum_{\nu=1}^{q-1} x^{q(q-1)+\nu}.$$

Lemma 11. *If $q \geq 2$, there exists x_q between 0 and 1 such that*

$$(59) \quad f_q(x_q) = \max_{0 \leq x \leq 1} f_q(x).$$

Moreover, we have the asymptotic formula

$$(60) \quad \int_0^1 f_q(x)^m x^{q-1} dx = f_q(x_q)^{m(1+o(1))} \quad \text{as } m \rightarrow \infty$$

and the bounds

$$(61) \quad \frac{1}{(q+1)e} < f_q(x_q) < \frac{1}{2e}.$$

Proof. Since

$$(62) \quad \frac{f'_q(x)}{x^{q(q-1)-1}} = q(q-1)^2 - \sum_{\nu=1}^{q-1} (q(q-1) + \nu)x^\nu$$

is decreasing for $x \geq 0$, positive at $x = 0$ and negative at $x = 1$, it follows that $f'_q(x)$ vanishes at exactly one point $x = x_q$ in the open interval $(0, 1)$. Then (59) holds with this choice of x_q , since $f_q(0) = f_q(1) = 0$ and $f_q(x) > 0$ for $0 < x < 1$. Differentiating (62), we substitute $f'_q(x_q) = 0$ and find that $f''_q(x_q) < 0$; by Laplace's method (see, e.g., [5], Section 4.2) we obtain (60).

Since

$$q - F_q(x) = (q - 1) \left(1 - \frac{x + x^2 + \cdots + x^{q-1}}{q - 1} \right),$$

the inequalities

$$(x \cdot x^2 \cdots x^{q-1})^{1/(q-1)} \leq \frac{x + x^2 + \cdots + x^{q-1}}{q - 1} \leq x \quad \text{for } 0 \leq x \leq 1$$

yield

$$(q - 1)(1 - x)x^{q(q-1)} \leq f_q(x) \leq (q - 1)(1 - x^{q/2})x^{q(q-1)}.$$

Using the relations

$$\frac{1}{(r + 1)e} < \max_{0 \leq y \leq 1} ((1 - y)y^r) = \frac{r^r}{(r + 1)^{r+1}} < \frac{1}{re} \quad \text{for } r > 0,$$

we set $y = x$, $r = q(q - 1)$ and deduce the lower bound in (61), then set $y = x^{q/2}$, $r = 2(q - 1)$ and obtain the upper bound. $\square$

The next result with $q = 2$ is comparable to Lemma 7 with $m = 2^n - 1$.

Lemma 12. *For $q \geq 2$, we have the bounds*

$$(63) \quad 0 < I'_{n,m,q} < \frac{q}{(2e)^m} \quad \text{with } m = \frac{q^n - 1}{q - 1}$$

and

$$(64) \quad -1 - \log(q + 1) < \lim_{n \rightarrow \infty} \frac{\log I'_{n,(q^n-1)/(q-1),q}}{(q^n - 1)/(q - 1)} < -1 - \log 2.$$

Proof. Let $m = (q^n - 1)/(q - 1)$, so that $x^{q^{n+1}-1} = x^{(mq+1)(q-1)}$. Using (53), (59) and Ramanujan's formula $I'_{0,0,q} = \gamma$, we have

$$\begin{aligned} (65) \quad 0 < I'_{n,m,q} &= \int_0^1 \frac{G_q(x)}{F_q(x)} (q - F_q(x))^m x^{(mq+1)(q-1)} \sum_{\nu=n+1}^{\infty} x^{q^\nu - q^{n+1}} dx \\ &< \int_0^1 f_q(x)^m \frac{G_q(x)}{F_q(x)} \left(1 + \sum_{\nu=1}^{\infty} x^{q^\nu - 1} \right) dx \\ &\leq f_q(x_q)^m (c_q + \gamma), \quad \text{where } c_q = \int_0^1 \frac{G_q(x)}{F_q(x)} dx, \end{aligned}$$

and (63) follows using $c_q + \gamma < (q - 1) + \gamma < q$ and the upper bound in (61).

We also have

$$(66) \quad I'_{n,m,q} > \int_0^1 \frac{G_q(x)}{F_q(x)} (q - F_q(x))^m x^{(mq+1)(q-1)} dx > \frac{q-1}{2} \int_0^1 f_q(x)^m x^{q-1} dx.$$

From (65), (66), (60) we deduce that

$$(67) \quad \lim_{n \rightarrow \infty} \frac{\log I'_{n,(q^n-1)/(q-1),q}}{(q^n - 1)/(q - 1)} = \log f_q(x_q),$$

and (61) gives (64). $\square$

Remark. The exact value of the constant c_q is $-\psi(1/q) - \log q - \gamma$, as another formula of Ramanujan shows (see [2], Proposition 2.1).

Theorem 9 (Base q Asymptotic Formula for γ). *If $q \geq 2$ and $m = (q^n - 1)/(q - 1)$, then*

$$\gamma = \frac{A'_{n,m,q} - L'_{n,m,q}}{q^m} + \delta_{m,q} \quad \text{with} \quad 0 < \delta_{m,q} < \frac{q}{(2eq)^m}.$$

Proof. Set $\delta_{m,q} = I'_{n,m,q}/q^m$ and use (58) and (63). $\square$

Remark. For $q = 2$, one could use $I'_{m,n,q}$ with $m = (q^n - 1)/(q - 1)$ to derive a rationality criterion and an irrationality test for γ (but they are already contained in Theorems 2 and 3 as the case $m = 2^n - 1$, since $I'_{n,2^n-1,2} = I_{n,2^n-1}$ and $L'_{n,2^n-1,2} = L_{n,2^n-1}$). For $q > 2$, this is not possible, the required inequality $d_{q^n} I'_{n,m,q} < 1$ not being valid, since

$$\lim_{n \rightarrow \infty} \frac{\log d_{q^n}}{q^n} = 1$$

(from the Prime Number Theorem) and the following observation imply that $d_{q^n} I'_{n,m,q}$ tends to infinity with n .

Lemma 13. *If $q \geq 3$, then*

$$(68) \quad \lim_{n \rightarrow \infty} \frac{\log I'_{n,(q^n-1)/(q-1),q}}{q^n} > -1.$$

Proof. By (64), we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\log I'_{n,(q^n-1)/(q-1),q}}{q^n} &= \frac{1}{q-1} \lim_{n \rightarrow \infty} \frac{\log I'_{n,(q^n-1)/(q-1),q}}{(q^n - 1)/(q - 1)} \\ &> -\frac{1 + \log(q+1)}{q-1} \quad \text{for } q \geq 2. \end{aligned}$$

Since the lower bound is increasing, and exceeds -1 when $q = 4$, this proves (68) for $q > 3$. For $q = 3$, we use (67) to replace the bound by the exact value

$$\frac{1}{2} \log f_3(x_3) = -0.90997390 \dots > -1,$$

where $x_3 = 0.86304075 \dots$ is the positive root of $12 - 7x - 8x^2$, which is the polynomial (62) with $q = 3$. $\square$

Remark. Alternatively, for q sufficiently large (68) follows from (67) and the formula

$$\lim_{q \rightarrow \infty} f_q(x_q) = \frac{1}{2e},$$

which in turn follows from the upper bound in (61) and the fact that the limit, as $q \rightarrow \infty$, of the lower bound in

$$\max_{0 \leq x \leq 1} f_q(x) \geq f_q(x) \Big|_{x=(1-1/q)^{1/q}} = \left(q - \frac{1/q}{1 - (1 - 1/q)^{1/q}} \right) \cdot \left(1 - \frac{1}{q} \right)^{q-1}$$

is $1/(2e)$.

10. CONCLUDING REMARKS

Any polynomial $P_m(x) \in \mathbb{Z}[x]$ can be used to generalize integral (18) to base q as follows:

$$(69) \quad \int_0^1 \left(\frac{q}{1-x^q} - \frac{1}{1-x} \right) P_m(x) \sum_{\nu=n+1}^{\infty} x^{q^\nu-1} dx.$$

Integrals (37) and (40) are particular cases of construction (69). The choice of $P_m(x)$, the “damping factor,” is based on the requirements:

- (i) to ensure the appearance of γ , and no other $\gamma_{j,q}$ from (39), in the corresponding linear forms;
- (ii) to produce sufficiently good asymptotics (upper estimates).

The choice $P_m(x) = (1-x)^m$ gives the best possible answer (38) to feature (ii); that is why we use it in Sections 2–4 for $q = 2$, and Section 8 for $q = 3$. The following lemma and the arguments in Section 7 show that $(1-x)^m$ cannot be used for $q > 3$, because of requirement (i).

Lemma 14. *If $q > 3$, the equality*

$$(70) \quad (1-x)^m = C + Q(x) \cdot \frac{1-x^q}{1-x}$$

is impossible for any $m > 0$, constant C and polynomial $Q(x)$.

Proof. For $q > 1$, equality (70) implies the system of equalities $(1 - \epsilon^j)^m = C$ for $j = 1, 2, \dots, q-1$, where ϵ is a primitive q -th root of unity. In particular, the value of $|1 - \epsilon^j|$ is the same for all $j = 1, 2, \dots, q-1$, which is possible only when $q = 2$ or $q = 3$. $\square$

One can avoid this difficulty by introducing linear combinations of powers of $1 - x$. More precisely, there exist (integer) constants $c_{m,0}, c_{m,1}, \dots, c_{m,q-1}$ such that

$$c_{m,0}(1-x)^m + c_{m,1}(1-x)^{m-1} + \dots + c_{m,q-1}(1-x)^{m-q+1} = C + Q(x) \cdot \frac{1-x^q}{1-x}.$$

However, we cannot choose the left-hand side as the polynomial $P_m(x)$, because the constants grow so much faster than m that requirement (ii) fails.

REFERENCES

1. W.N. Bailey, *Generalized Hypergeometric Series*, Cambridge Math. Tracts, vol. 32, Cambridge Univ. Press, Cambridge, 1935; 2nd reprinted edition, Stechert-Hafner, New York–London, 1964.
2. B.C. Berndt and D.C. Bowman, *Ramanujan's short unpublished manuscript on integrals and series related to Euler's constant*, Constructive, Experimental, and Nonlinear Analysis (Limoges, 1999) (M. Thera, ed.), CMS Conf. Proc., vol. 27, Amer. Math. Soc., Providence, RI, 2000, pp. 19–27.
3. P. Borwein, *On the irrationality of $\sum \frac{1}{q^n+r}$* , J. Number Theory **37** (1991), 253–259.
4. T. Bromwich, *An Introduction to the Theory of Infinite Series*, 2nd edition, Macmillan, London, 1926.
5. N.G. de Bruijn, *Asymptotic Methods in Analysis*, Dover Publications, New York, 1981.
6. R.W. Gosper, Jr., *Personal communication* (7 May 2002).
7. J.B. Rosser and L. Schoenfeld, *Approximate formulas for some functions of prime numbers*, Illinois J. Math. **6** (1962), 64–94.
8. P. Sebah, *Personal communication* (17 December 2002).
9. J. Sondow, *Criteria for irrationality of Euler's constant*, Proc. Amer. Math. Soc. (to appear).
10. J. Sondow, *A hypergeometric approach, via linear forms involving logarithms, to irrationality criteria for Euler's constant*, CRM Conference Proceedings of CNTA 7 (May 2002) (submitted).
11. G. Vacca, *A new series for the Eulerian constant $\gamma = .577\dots$* , Quart. J. Pure Appl. Math. **41** (1909–10), 363–364.

209 WEST 97TH STREET

NEW YORK, NY 10025 USA

URL: <http://home.earthlink.net/~jsondow/>

E-mail address: jsondow@alumni.princeton.edu

MOSCOW LOMONOSOV STATE UNIVERSITY

DEPARTMENT OF MECHANICS AND MATHEMATICS

VOROBIOVY GORY, GSP-2, MOSCOW 119992 RUSSIA

URL: <http://wain.mi.ras.ru/index.html>

E-mail address: wadim@ips.ras.ru